



DigestZyme

DigestZyme begins the process of food digestion in the stomach, reducing digestive stress on the pancreas.

ASSISTS COMPLETE AND EFFICIENT FOOD DIGESTION!

SUPPLEMENT FACTS		
Serving Size 2 capsules		
Amount Per Serving		% Daily Value
Enzyme proprietary blend	412 mg	
Amylase	12,200 DU	*
Protease	66,500 HUT	*
Lipase	400 FCCLU	*
Cellulase	1320 CU	*
Invertase	0.4 IAU	*
Malt Diastase	500 DP*	*
Lactase	800 ALU	*
Lactobacillus acidophilus	10 mg	*
Bifidobacterium longum	3 mg	*
*Daily Value not established		

Other ingredients: Vegetarian Capsule (cellulose & water)
Enzyme activity is measured in Food Chemical Codex (FCC) units.
Store tightly in a cool, dry place. Keep out of reach of children

120 capsules per bottle. No fillers.

Recommended Usage: Take 2 or more capsules with every meal. Take 1 - 2 capsules with snacks. The dosage is dependent on both the types of foods eaten and the individual's digestive difficulties. If alcohol is consumed with the meal, then the dosage should be increased. (1 extra capsule is suggested with wine.)

Indications:

- Most digestive disorders
- Dairy intolerance
- Gallbladder stress (see below)
- Prevention of malabsorption
- Intestinal toxemia
- Food allergies
- Indigestion
- Sugar intolerance

Gallbladder stress may show itself with these indications:

- Nausea following meal (anywhere from directly after to several hours later)
- Headache prior to or following meal
- Nausea that is relieved by eating or with a bowel movement
- Diarrhea following meals or thereafter
- Pain or tenderness in the upper abdomen

- Diarrhea or constipation
- Burps or belches
- Bloating
- A feeling of tenderness where the rib cage comes together

DIGESTZYME should be used for any of the symptoms mentioned above. Take 2-3 capsules with meals and, as needed, 1-3 capsules any other time. Symptoms should dissipate within fifteen to thirty minutes. Some people may require an additional 3 capsules of GASTROZYME for diarrhea or heartburn.

Individuals with extreme gallbladder or gastrointestinal problems may benefit from a regimen of GASTROZYME for one week, followed by a continuing program of DIGESTZYME along with GASTROZYME.

Contraindications:

In cases of Gastritis (extreme case), Gastric Ulcers, or Duodenal Ulcers, use GASTROZYME with meals and/or as needed.

Although DIGESTZYME is available in a powdered form, the capsules can also be pulled apart and may then be used for children, pets, or in cases where a person is unable to swallow capsules. Powdered DigestZyme should be mixed with acceptable food in a teaspoon for a child or with baby formula in a spoon feeder for a baby. Do not add DIGESTZYME to a bottle of baby formula or to a bowl of food, since it will automatically digest the food in that container.